



LISBON
ADDICTIONS
2019



UNIDADE
RECOMEÇO
HELVETIA



Crackland in Brazil: Epidemiological Monitoring Method and Results

Authors:

Clarice Sandi Madruga, PhD;
Gleuda Apolinário,
Claudio Jerônimo, PhD;
Ronaldo Laranjeira, MD, PhD

Presenter:

Clarice S Madruga*

The authors declare no conflict of interest.

* Graduated in Psychology, Masters degree in Neurosciences and in Substance Misuse (Sussex University). Doctorate degree in Psychiatry and Medical Psychology (UNIFESP and Kings College London). Affiliated professor at UNIFESP's Department of Psychiatry. Consultor for the State of Sao Paulo's Council of Drug Policy via the United Nations Development Programme.



LISBON
ADDICTIONS
2019

Crack Use in Brazil:



II BNADS
Brazilian National Alcohol
and Drugs Survey

Addictive Behaviors 39 (2014) 297–301



Contents lists available at ScienceDirect

Addictive Behaviors



Prevalence of Cocaine Use in Brazil: Data from the II Brazilian National Alcohol and Drugs Survey (BNADS)

Renata Rigacci Abdalla^{a,*}, Clarice S. Madruga^a, Marcelo Ribeiro^a, Ilana Pinsky^a,
Raul Caetano^b, Ronaldo Laranjeira^a

^a National Institute of Policies on Alcohol and Drugs (INPAD) of the Psychiatry Department of the Federal University of São Paulo (UNIFESP), Brazil

^b University of Texas School of Public Health, Dallas Regional Campus, USA



HIGHLIGHTS

- Crack/cocaine last year consumption rate was 2.2% (3.7% in men and 0.7% in women)
- Snorted cocaine last year consumption rate was 1.7% (2.9% in men and 0.6% in women)
- Smoked cocaine last year consumption rate was 0.8% in Brazil
- Cocaine addiction prevalence was 0.6% in the population and 41.4% among users
- Younger participants and men were more likely to use cocaine.

Crackland







Method


Time Location sampling:



Occupancy Area 2016




Occupancy Area 2017 - 1







Occupancy Area 2017-2





Occupancy Area 2019










Likert scale motivational stages



I don't want...

I really want...

44.6%

- Declared really want to quit

Motivation associated factors

- History of previous treatments was significantly associated with motivation to quit

Our Team

Coordenação da pesquisa:

Clarice S Madruga, Phd

Professora Pós-Graduação em Psiquiatria e Psicologia Médica, UNIFESP

Email: clarice.madruga@unifesp.br

Entrevistadores 2016:

- Paulo Roberto da Silva

Conselheiro Tenda Recomeço, Especialista em Dependência Química UNIAD/UNIFESP

- Lays dos Santos Rodrigues

Assistente social, Especialista em Dependência Química UNIAD/UNIFESP

- Mônica Rabello

Psicóloga, Especialista em Dependência Química UNIAD/UNIFESP

2017:

- Patrícia Bernardete

Psicóloga, Especialista em Dependência Química UNIAD/UNIFESP, Mestranda Psiquiatria UNIFESP

- Tainá Saboia

Tecnologia da informação

