The Association between Moral Injury and Cannabis Use Disorder among Israeli Combat Veterans

Daniel Feingold, PhD

Moral Injury

The event

- o Transgression by **self**
- o Transgression by **others**
- Betrayal

Long-term consequences

- Emotional:
 - Anger, disgust, self-condemnation
- Interpersonal: Isolation, exclusion
- Behavioral:

Self-handicapping, self harm, <u>Substance use</u>




Contents lists available at ScienceDirect Clinical Psychology Review



Moral injury and moral repair in war veterans: A preliminary model and intervention strategy

Brett T. Litz $^{\rm a,b,*}$, Nathan Stein $^{\rm a}$, Eileen Delaney $^{\rm a}$, Leslie Lebowitz $^{\rm c}$, William P. Nash $^{\rm c}$, Caroline Silva $^{\rm a}$, Shira Maguen $^{\rm d}$

Shame & Guilt






Past-year cannabis use 204,000,000 Percentage of population aged 15-64 8.00 5.01 - 8.00 4.01 - 6.00 2.01 - 4.00 <=2.00 No data provided Data older than 2006

Note: The boundaries and names shown and the designations used on this map do not imply official endorsement or acceptance by the United Nations. Dashed lines represent undetermined boundaries. Dotted line represents approximately the Line of Control in Jammu and Kashmir agreed upon by India and Pakistan. The final status of Jammu and Kashmir has not yet beenagreed upon by the parties. Final boundary between the Republic of South South nas not yet been determined.

% reporting past-year use of cannabis



International Journal of Mental Health and Addiction (2019) 17:217–233 https://doi.org/10.1007/s11469-018-0012-8

ORIGINAL ARTICLE



The Association Between Moral Injury and Substance Use Among Israeli Combat Veterans: the Mediating Role of Distress and Perceived Social Support

Daniel Feingold¹ · Gadi Zerach¹ · Yossi Levi-Belz^{2,3}

Participants: 191 combat veterans


Measurements:

• Moral Injury Event Scale (MIES) Nash et al., 2013.


e.g., "I acted in ways that violated my own moral code or values"


- Patient Health Questionnaire (PHQ-8) Kroenke et al. 2009
- Posttraumatic Stress Disorder Checklist (PCL-5) Bovin et al. 2016
- Multidimensional Scale of Perceived Social Support Zimet 1988
- Past-year frequency of various substance use

% reporting past-year use of cannabis



Results





Feingold et al. (2020). DAR; Koob et al. (2020). Am J Psychiatry



Aviya Ashwal-Malka^a, Keren Tal-Kishner^b, Daniel Feingold^{a,*}

Participants: 215 male combat veterans ≥ 1 year of combat service regular cannabis use

Instruments:

Cannabis Use Disorder Identification Test Revised (CUDIT-R; Adamson et al., 2010)


Moral Injury Event Scale (MIES; Nash et al., 2013)

Patient Health Questionnaire Depression Scale (PHQ-8; Kroenke et al., 2009)

The Multidimensional Scale of Perceived Social Support (MSPSS; Zimet et al., 1988)

Results


o Positive association between moral injury and cannabis CUD symptoms o Mediated by depression




Results

Social support moderates the association between moral injury and cannabis CUD






MIES - betrayal





"Only time heals a wounded heart..." Araybia S. L. Williams,





MI - betrayal

Lewenstein et al. (in preparation)

CUD symptoms

Narratives of young veterans who use cannabis

From Enchantment to Disillusion: A Narrative Exploration of Cannabis Use Disorder Among Young Israeli Combat Veterans

Miri Serebro¹, Shira Sobol-Goldberg² and Daniel Feingold^{3*}

- Combat veterans ≤ 5 years since release from duty (n=12).
- Regular use of cannabis.
- In-depth interviews.
- Systematic content analysis

Narratives of young veterans who use cannabis

From Enchantment to Disillusion: A Narrative Exploration of Cannabis Use Disorder Among Young Israeli Combat Veterans

Miri Serebro¹, Shira Sobol-Goldberg² and Daniel Feingold³⁺





"I made it through this war, I felt I deserve to do whatever I want".

"I'm starting to feel bad about weed. When you're weak you start to see its downsides."


Narratives of older veterans who use cannabis



- Combat veterans >5 years since release from duty (n=10)
- Regular use of cannabis
- In-depth interviews
- Systematic content analysis

Narratives of older veterans who use cannabis







- Moral injury is closely associated with cannabis use and CUD.
- This association may be mediated by emotional distress.
- Paradoxically, social support may elevate the MI-cannabis correlation.
- Time and finding meaning may increase resilience.

THANK YOU for LISTENING

Daniel Feingold, PhD danielfe@ariel.ac.il